

CHILEAN JOURNAL OF STATISTICS

Edited by Víctor Leiva and Carolina Marchant

A free open access journal indexed by



Web of
Science
Group

Scopus®



ELSEVIER

Volume 11 Number 2
December 2020

ISSN: 0718-7912 (print)

ISSN: 0718-7920 (online)

Published by the
Chilean Statistical Society

SOCH E 
SOCIEDAD CHILENA DE ESTADÍSTICA

AIMS

The Chilean Journal of Statistics (ChJS) is an official publication of the Chilean Statistical Society (www.soche.cl). The ChJS takes the place of *Revista de la Sociedad Chilena de Estadística*, which was published from 1984 to 2000.

The ChJS covers a broad range of topics in statistics, as well as in artificial intelligence, big data, data science, and machine learning, focused mainly on research articles. However, review, survey, and teaching papers, as well as material for statistical discussion, could be also published exceptionally. Each paper published in the ChJS must consider, in addition to its theoretical and/or methodological novelty, simulations for validating its novel theoretical and/or methodological proposal, as well as an illustration/application with real data.

The ChJS editorial board plans to publish one volume per year, with two issues in each volume. On some occasions, certain events or topics may be published in one or more special issues prepared by a guest editor.

EDITORS-IN-CHIEF

Víctor Leiva
Carolina Marchant

Pontificia Universidad Católica de Valparaíso, Chile
Universidad Católica del Maule, Chile

EDITORS

Héctor Allende Cid	<i>Pontificia Universidad Católica de Valparaíso, Chile</i>
Danilo Alvares	<i>Pontificia Universidad Católica de Chile</i>
José M. Angulo	<i>Universidad de Granada, Spain</i>
Robert G. Aykroyd	<i>University of Leeds, UK</i>
Narayanaswamy Balakrishnan	<i>McMaster University, Canada</i>
Michelli Barros	<i>Universidade Federal de Campina Grande, Brazil</i>
Carmen Batanero	<i>Universidad de Granada, Spain</i>
Ionut Bebu	<i>The George Washington University, US</i>
Marcelo Bourguignon	<i>Universidade Federal do Rio Grande do Norte, Brazil</i>
Márcia Branco	<i>Universidade de São Paulo, Brazil</i>
Oscar Bustos	<i>Universidad Nacional de Córdoba, Argentina</i>
Luis M. Castro	<i>Pontificia Universidad Católica de Chile</i>
George Christakos	<i>San Diego State University, US</i>
Enrico Colosimo	<i>Universidade Federal de Minas Gerais, Brazil</i>
Gauss Cordeiro	<i>Universidade Federal de Pernambuco, Brazil</i>
Francisco Cribari-Neto	<i>Universidade Federal de Pernambuco, Brazil</i>
Francisco Cysneiros	<i>Universidade Federal de Pernambuco, Brazil</i>
Mário de Castro	<i>Universidade de São Paulo, São Carlos, Brazil</i>
José A. Díaz-García	<i>Universidad Autónoma Agraria Antonio Narro, Mexico</i>
Raul Fierro	<i>Universidad de Valparaíso, Chile</i>
Jorge Figueroa-Zúñiga	<i>Universidad de Concepción, Chile</i>
Isabel Fraga	<i>Universidade de Lisboa, Portugal</i>
Manuel Galea	<i>Pontificia Universidad Católica de Chile</i>
Diego Gallardo	<i>Universidad de Atacama, Chile</i>
Christian Genest	<i>McGill University, Canada</i>
Viviana Giampaoli	<i>Universidade de São Paulo, Brazil</i>
Marc G. Genton	<i>King Abdullah University of Science and Technology, Saudi Arabia</i>
Patricia Giménez	<i>Universidad Nacional de Mar del Plata, Argentina</i>
Hector Gómez	<i>Universidad de Antofagasta, Chile</i>
Yolanda Gómez	<i>Universidad de Atacama, Chile</i>
Emilio Gómez-Déniz	<i>Universidad de Las Palmas de Gran Canaria, Spain</i>
Daniel Griffith	<i>University of Texas at Dallas, US</i>
Eduardo Gutiérrez-Peña	<i>Universidad Nacional Autónoma de México</i>
Nikolai Kolev	<i>Universidade de São Paulo, Brazil</i>
Eduardo Lalla	<i>University of Twente, Netherlands</i>
Shuangzhe Liu	<i>University of Canberra, Australia</i>
Jesús López-Fidalgo	<i>Universidad de Navarra, Spain</i>
Liliana López-Kleine	<i>Universidad Nacional de Colombia</i>
Rosângela H. Loschi	<i>Universidade Federal de Minas Gerais, Brazil</i>
Manuel Mendoza	<i>Instituto Tecnológico Autónomo de México</i>
Orietta Nicolis	<i>Universidad Andrés Bello, Chile</i>
Ana B. Nieto	<i>Universidad de Salamanca, Spain</i>
Teresa Oliveira	<i>Universidade Aberta, Portugal</i>
Felipe Osorio	<i>Universidad Técnica Federico Santa María, Chile</i>
Carlos D. Paulino	<i>Instituto Superior Técnico, Portugal</i>
Fernando Quintana	<i>Pontificia Universidad Católica de Chile</i>
Nalini Ravishanker	<i>University of Connecticut, US</i>
Fabrizio Ruggeri	<i>Consiglio Nazionale delle Ricerche, Italy</i>
José M. Sarabia	<i>Universidad de Cantabria, Spain</i>
Helton Saulo	<i>Universidade de Brasília, Brazil</i>
Pranab K. Sen	<i>University of North Carolina at Chapel Hill, US</i>
Julio Singer	<i>Universidade de São Paulo, Brazil</i>
Milan Stehlik	<i>Johannes Kepler University, Austria</i>
Alejandra Tapia	<i>Universidad Católica del Maule, Chile</i>
M. Dolores Ugarte	<i>Universidad Pública de Navarra, Spain</i>

CONTENTS

Víctor Leiva and Carolina Marchant <i>Confirming our international presence with publications and submissions from all continents in COVID-19 pandemic</i>	69
Ibrahim M. Almanjahie, Mohammed Kadi Attouch, Omar Fetitah, and Hayat Louhab <i>Robust kernel regression estimator of the scale parameter for functional ergodic data with applications</i>	73
Ricardo Puziol de Oliveira, Marcos Vinicius de Oliveira Peres, Jorge Alberto Achcar, and Nasser Davarzani <i>Inference for the trivariate Marshall-Olkin-Weibull distribution in presence of right-censored data</i>	95
Henrique José de Paula Alves and Daniel Furtado Ferreira <i>On new robust tests for the multivariate normal mean vector with high-dimensional data and applications</i>	117
Josmar Mazucheli, André F.B. Menezes, Sanku Dey, and Saralees Nadarajah <i>Improved parameter estimation of the Chaudhry and Ahmad distribution with climate applications</i>	137
André Leite, Abel Borges, Geiza Silva, and Raydonal Ospina <i>A timetabling system for scheduling courses of statistics and data science: Methodology and case study</i>	151
Jorge Figueroa-Zúñiga, Rodrigo Sanhueza-Parkes, Bernardo Lagos-Álvarez, and Germán Ibacache-Pulgar <i>Modeling bounded data with the trapezoidal Kumaraswamy distribution and applications to education and engineering</i>	163

STATISTICAL INFERENCE
RESEARCH PAPER

Improved parameter estimation of the Chaudhry and Ahmad distribution with climate applications

JOSMAR MAZUCHELI^{1,*}, ANDRÉ F.B. MENEZES², SANKU DEY³, and SARALEES NADARAJAH⁴

¹Departamento de Estatística, Universidade Estadual de Maringá, Maringá-PR, Brazil,

²Departamento de Estatística, Universidade Estadual de Campinas, Campinas-SP, Brazil,

³Department of Statistics, St. Anthony's College, Shillong, Meghalaya, India

⁴School of Mathematics, University of Manchester, Manchester M13 9PL, UK

(Received: 04 August 2020 · Accepted in final form: 11 December 2020)

Abstract

The Chaudhry-Ahmad distribution is a two-parameter continuous probability distribution obtained as a solution to a generalized Pearson system of differential equation. Although its probability density curve resembles the inverse-Gaussian, gamma, log-normal, Weibull and other distributions, it has been neglected in the analysis of right-skewed data. The purpose of this paper is three folded. Firstly, to reparametrize the Chaudhry and Ahmad distribution and present some of its basic properties. Secondly to derive the analytical bias-corrected maximum likelihood estimators applying the Cox-Snell methodology and thirdly to study, by MC simulations, the small-sample properties of the maximum likelihood estimators and their bias-corrected versions, obtained from the Cox-Snell formula and by parametric bootstrap technique. The numerical results show, for both parameters, that the maximum likelihood estimators are highly biased, especially in small samples. On the other hand, both, the analytical and bootstrap methodologies, significantly reduce the biases and the mean-squared errors. It is apparent from the results that the analytical bias-correction is more efficient than bootstrap resamples. Finally, wind speed data from six weather stations distributed in the state of Tocantins in Brazil is used to illustrate the applicability of the proposed methods.

Keywords: Bootstrap bias correction · Cox-Snell bias-correction · Maximum likelihood estimation · Monte Carlo simulation · Wind speed data.

Mathematics Subject Classification: Primary 60E05 · Secondary 62F10.

1. INTRODUCTION

Chaudhry and Ahmad (1993) introduced a nonnegative two-parameter probability distribution, called the Chaudhry-Ahmad (CA) distribution as a solution of the generalized Pearson system of differential equation. It is noteworthy that from the generalized Pearson system of probability distributions, many continuous probability density functions (PDFs) can be generated (Sankaran et al., 2003; Stavroyiannis, 2014). Indeed, as discussed in Shakil et al. (2010, 2016), the well known families of distributions such as the normal and

*Corresponding author. Email: jmazucheli@gmail.com

Student-t (known as Pearson type VII), beta distribution (known as Pearson type I) and gamma distribution (known as Pearson type III), introduced by Karl Pearson during the late 19th century (Pearson, 1893, 1895, 1901, 1916), can be generated as a solution to Equation (1) by proper choice of its parameters.

Although the CA PDF curve resembles the inverse-Gaussian (IG), gamma, log-normal, Weibull and other distributions, it has not been widely explored in the statistical literature. Recently, Shakil et al. (2010) derived a family of distribution, which includes the CA distribution as a special case. To the best of our knowledge, there are only two real data analysis considering the CA distribution. Nanos and Montero (2001) showed that CA distribution fitted better than the Weibull distribution in a problem involving prediction of the diameter distribution of a stand. In Nanos et al. (2000) the Weibull and CA distributions were used to model resin production distributions for maritime pine stands.

It is important to point out that the CA distribution is capable of modeling increasing hazard rate functions (HRFs). There are many situations where only increasing HRFs are used or observed: Woosley and Cossman (2007) observed that drugs during clinical development have increasing HRFs; Tsarouhas and Arvanitoyannis (2010) showed that machines of the bread production display increasing HRFs; Koutras (2011) observed that software degradation times have increasing HRFs; Lai (2013) investigated the optimum number of minimal repairs for systems have increasing hazard rates and so on.

Although the maximum likelihood (ML) estimators have many appealing properties (Edwards, 1992; Lehmann and Casella, 1998), it is also well known that ML estimators could be biased, especially when the study is being done in small samples. Owing to this reason, researchers strive to develop nearly unbiased estimators for the parameters of several probability distributions. Notable among them are Saha and Paul (2005), Lemonte et al. (2007), Giles and Feng (2009), Lagos-Álvarez et al. (2011), Giles (2012a), Giles (2012b), Schwartz et al. (2013), Giles et al. (2013), Teimouri and Nadarajah (2013), Ling and Giles (2014), Zhang and Liu (2015), Teimouri and Nadarajah (2016), Reath (2016), Schwartz and Giles (2016), Wang and Wang (2017), Mazucheli and Dey (2018), Mazucheli et al. (2018), Mazucheli et al. (2020) and references cited therein.

The objective of this paper is to perform improved parameter estimation of the CA distribution. We consider the analytical methodology introduced by Cox-Snell (1968) and the parametric bootstrap resampling method (Efron, 1982). We describe two corrective approaches to bias-correction, both methods reduce the biases of the ML estimators to the second order magnitude.

After this introduction, the paper is organized as follows. In Section 2, we introduce the CA distribution and deduce expressions used to obtain the ML estimators of its parameters, calculating the expected Fisher information matrix. In Section 3, by using the Cox-Snell formula, we derive analytical expressions for the second order biases of the maximum likelihood estimators, and also discuss the bootstrap bias correction. A Monte Carlo (MC) simulation study is carried out in Section 4 to compare the ML estimators and their bias-corrected versions, obtained from the Cox-Snell formula and parametric bootstrap technique. An application by using wind speed data from Brazil is provided also in this section. As a result of this application, we are able to provide, for example, better estimates of most frequent wind speeds observed at various stations. Some concluding remarks are presented in Section 5.

2. PRELIMINARIES, MODEL DESCRIPTION AND ESTIMATION

In this section, we provide background on the CA distribution and the ML estimators of its parameters, as well as the corresponding expected Fisher information matrix.

2.1 BACKGROUND ON THE CHAUDHRY-AHMAD DISTRIBUTION

Chaudhry and Ahmad (1993) developed a two-parameter probability distribution as a solution of the generalized Pearson system of differential equation

$$\frac{d}{dx} f(x) = \frac{c_0 + c_1 x + c_2 x^2 + \cdots + c_m x^m}{c'_0 + c'_1 x + c'_2 x^2 + \cdots + c'_n x^n} f(x), \quad (1)$$

where $m, n \geq 1$ are integers, and the coefficients c and c' are real numbers. These authors considered a special case of Equation (1) taking $m = 4$, $n = 3$, $c'_0 = c'_1 = c'_2 = 0$, $c_4/2c'_3 = -2\alpha$, $c_0/2c'_3 = 2\beta$ and $c'_3 \neq 0$. This distribution, which now bear their names, can also be obtained as the root reciprocal of the inverse Gaussian distribution, that is, the distribution of the random variable $X = 1/\sqrt{Y}$, where $Y \sim \text{IG}(\mu, \lambda)$ with $\mu = (\alpha/\beta)^{1/2}$ and $\lambda = 2\alpha$.

The cumulative distribution function (CDF) of the CA distribution is given by

$$F(x; \alpha, \beta) = \Phi \left[\sqrt{2} \left(\sqrt{\alpha} x - \sqrt{\beta} x^{-1} \right) \right] - \exp \left(4 \sqrt{\alpha \beta} \right) \Phi \left[-\sqrt{2} \left(\sqrt{\alpha} x + \sqrt{\beta} x^{-1} \right) \right], \quad (2)$$

where $x, \alpha, \beta > 0$ and Φ denotes the CDF of a standard normal distribution.

Solving the orthogonality differential equation of Cox and Reid (1987), we consider in Equation (2) $\beta = \alpha\lambda^4$, such that λ will be the mode of the PDF. The advantage of such parametrization is that λ has a direct interpretation and it is orthogonal to α . Thus, from Equation (2), the PDF of a CA distributed random variable with parameters α and λ can be written as

$$f(x; \alpha, \lambda) = 2 \sqrt{\frac{\alpha}{\pi}} \exp \left[- \left(\sqrt{\alpha} x - \lambda^2 \sqrt{\alpha} x^{-1} \right)^2 \right]. \quad (3)$$

Figure 1 displays the PDF and the HRF curves considering different values of α and $\lambda = 1$ (λ is a location parameter). We observe that the PDF is skewed to the right and unimodal with turning point at $x_{\max} = \lambda = 1$. We also observe that the HRF of CA distribution is monotone increasing.




Figure 1. PDF and HRF of the CA distribution for $\alpha = (0.5, 1.0, 2.0 \text{ and } 4.0)$ and $\lambda = 1$.

The k th moment about the origin of CA distribution is given by

$$\mu'_k = 2 \sqrt{\frac{\alpha}{\pi}} \exp(2\alpha\lambda^2) \lambda^{k+1} K_{\frac{k}{2}+\frac{1}{2}}(2\alpha\lambda^2), \quad (4)$$

where K_ν denotes the modified Bessel function of the second kind (Abramowitz and Stegun, 1974). In particular, from Equation (4), the first four moments about the origin are stated as

$$\begin{aligned} \mu'_1 &= 2 \sqrt{\frac{\alpha}{\pi}} \exp(2\alpha\lambda^2) \lambda^2 K_{\frac{1}{2}+\frac{1}{2}}(2\alpha\lambda^2), \\ \mu'_2 &= \frac{2\alpha\lambda^2 + 1}{2\alpha}, \\ \mu'_3 &= 2 \sqrt{\frac{\alpha}{\pi}} \exp(2\alpha\lambda^2) \lambda^4 K_{\frac{3}{2}+\frac{1}{2}}(2\alpha\lambda^2), \\ \mu'_4 &= \frac{4\alpha^2\lambda^4 + 6\alpha\lambda^2 + 3}{4\alpha^4}. \end{aligned}$$

2.2 MAXIMUM LIKELIHOOD ESTIMATION

Suppose that $\mathbf{X} = (X_1, \dots, X_n)^\top$ is a random sample of size n from CA distribution with PDF given by Equation (3) and $\mathbf{x} = (x_1, \dots, x_n)^\top$ its observations. The log-likelihood function for $\boldsymbol{\theta} = (\alpha, \lambda)$ is given by

$$\ell(\boldsymbol{\theta}; \mathbf{x}) \propto \frac{n}{2} \log(\alpha) + 2n\alpha\lambda^2 - \alpha \sum_{i=1}^n x_i^2 - \lambda^4 \alpha \sum_{i=1}^n x_i^{-2}. \quad (5)$$

Differentiating in Equation (5) with respect to α and λ , we have the score vector $\mathbf{U}_{\boldsymbol{\theta}} = (U_\alpha, U_\lambda)^\top$ with components given by

$$U_\alpha = \frac{n}{2\alpha} + 2n\lambda^2 - \sum_{i=1}^n x_i^2 - \lambda^4 \sum_{i=1}^n x_i^{-2}, \quad (6)$$

$$U_\lambda = 4n\alpha\lambda - 4\lambda^3\alpha \sum_{i=1}^n x_i^{-2}. \quad (7)$$

After simple algebraic manipulation of Equations (6) and (7), note that the ML estimates of α and λ can be written as $\hat{\lambda} = (m'_{-2})^{-1/2}$ and $\hat{\alpha} = [2(m'_{-2}m'_2 - 1)m'_{-2}]^{-1}(m'_{-2})^2$, where $m'_2 = (1/n) \sum_{i=1}^n x_i^2$ and $m'_{-2} = (1/n) \sum_{i=1}^n x_i^{-2}$.

The expected Fisher information matrix of $\boldsymbol{\theta}$ is given by

$$\mathbf{I}(\boldsymbol{\theta}) = [I_{ij}] = -nE \left(\frac{\partial^2}{\partial \theta_i \partial \theta_j} \log(f(x_i; \boldsymbol{\theta})) \right) = \begin{bmatrix} -\frac{n}{2\alpha^2} & 0 \\ 0 & -8n\alpha \end{bmatrix}, \quad i, j = 1, 2. \quad (8)$$

From Equation (8), we observe that the information matrix is diagonal, which means that the ML estimators are asymptotically independent. Hence, the asymptotic variance

of $\hat{\alpha}$ and $\hat{\lambda}$ are given, respectively, by

$$\text{Var}(\hat{\alpha}) = \frac{2\alpha^2}{n}, \quad \text{Var}(\hat{\lambda}) = \frac{1}{8n\alpha}. \quad (9)$$

The asymptotic variance of $\hat{\lambda}$ only depends on α . Thus, as α decreases, the variance of $\hat{\lambda}$ increases. The asymptotic $100(1 - \delta)$ confidence intervals for α and λ can be obtained respectively as

$$\hat{\alpha} \pm z_{\delta/2} \sqrt{\widehat{\text{Var}}(\hat{\alpha})}, \quad \hat{\lambda} \pm z_{\delta/2} \sqrt{\widehat{\text{Var}}(\hat{\lambda})}, \quad (10)$$

where $z_{\delta/2}$ indicated in Equation (10) denotes the $100(1 - \delta/2)$ percentile of the standard normal distribution.

3. BIAS-CORRECTED MAXIMUM LIKELIHOOD ESTIMATORS

In this section, we derive analytical expressions for the second order biases of the maximum likelihood estimators by using the Cox-Snell formula, and also discuss the bootstrap bias correction.

3.1 COX-SNELL ANALYTIC BIAS CORRECTION

Let $\ell(\boldsymbol{\theta}; \mathbf{x})$ denote the log-likelihood function of a p -dimensional parameter vector $\boldsymbol{\theta}$ based on a sample of observations \mathbf{x} . We assume the following regularity conditions on the behavior of the log-likelihood function (Cox and Hinkley, 1979):

- (a) X_i , for $i = 1, \dots, n$, are independent and identically distributed random variables.
- (b) The parameter space of $\boldsymbol{\theta}$ is compact.
- (c) The true but unknown parameter value $\boldsymbol{\theta}_0$ is identified, that is,

$$\boldsymbol{\theta}_0 = \arg \max_{\boldsymbol{\theta}} \text{E}_{\boldsymbol{\theta}_0} [\log (f(x_i; \boldsymbol{\theta}))].$$

- (d) The likelihood function

$$\ell(\boldsymbol{\theta}; \mathbf{x}) = \sum_{i=1}^n \log (f(x_i; \boldsymbol{\theta}))$$

is continuous in $\boldsymbol{\theta}$.

- (e) $\text{E}_{\boldsymbol{\theta}_0} [\log (f(x_i; \boldsymbol{\theta}))]$ exists.
- (f) The log-likelihood function is such that $(1/n)\ell(\boldsymbol{\theta}; \mathbf{x})$ converges almost surely (in probability) to $\text{E}_{\boldsymbol{\theta}_0} [\log(f(x_i; \boldsymbol{\theta}))]$ uniformly in $\boldsymbol{\theta}$.

Conditions (a) to (d) are clearly satisfied for the CA distribution. Conditions (e) and (f) are also satisfied since, for all $\alpha > 0$ and $\beta > 0$,

$$\int_0^\infty x^2 \exp \left(-\alpha x^2 - \frac{\lambda^4 \alpha}{x^2} \right) dx < \infty,$$

$$\int_0^\infty x^{-2} \exp \left(-\alpha x^2 - \frac{\lambda^4 \alpha}{x^2} \right) dx < \infty.$$

The joint cumulants of the derivatives of ℓ are given by

$$\mathbf{I}_{ij} = \mathbb{E} \left[\frac{\partial^2 \ell}{\partial \boldsymbol{\theta}_i \partial \boldsymbol{\theta}_j} \right], \quad \mathbf{I}_{ijl} = \mathbb{E} \left[\frac{\partial^3 \ell}{\partial \boldsymbol{\theta}_i \partial \boldsymbol{\theta}_j \partial \boldsymbol{\theta}_l} \right], \quad \mathbf{I}_{ij,l} = \mathbb{E} \left[\left(\frac{\partial^2 \ell}{\partial \boldsymbol{\theta}_i \partial \boldsymbol{\theta}_j} \right) \left(\frac{\partial \ell}{\partial \boldsymbol{\theta}_l} \right) \right],$$

for $i, j, l = 1, \dots, p$. All these expression are assumed to be of order $\mathcal{O}(n)$.

Cox-Snell (1968) showed that when the samples are independent, but not necessarily identically distributed, the bias of the r th element of the ML estimator of $\boldsymbol{\theta}$, $\hat{\boldsymbol{\theta}}$, can be expressed as

$$\mathcal{B}(\hat{\boldsymbol{\theta}}_r) = \sum_{i=1}^p \sum_{j=1}^p \sum_{l=1}^p \mathbf{I}^{ri} \mathbf{I}^{jl} [0.5 \mathbf{I}_{ijl} + \mathbf{I}_{ij,l}] + \mathcal{O}(n^{-2}), \quad (11)$$

where $r = 1, \dots, p$ and \mathbf{I}^{ij} denotes the (i, j) th element of the inverse of the expected Fisher information matrix.

In respect to the orthogonally parametrization of CA distribution, after extensive algebra, it can be shown that $\mathbf{I}_{111} = -24n\alpha/\lambda$, $\mathbf{I}_{122} = \mathbf{I}_{212} = \mathbf{I}_{221} = -8n$, $\mathbf{I}_{222} = n/\alpha^3$, $\mathbf{I}_{11,1} = 24n\alpha/\lambda$, $\mathbf{I}_{12,2} = \mathbf{I}_{21,2} = 8n$ and all other terms are equal to zero. Hence, the second-order bias of the ML estimators of α and λ are given respectively by

$$\mathcal{B}(\hat{\alpha}) = \frac{3\alpha}{n} \quad (12)$$

and

$$\mathcal{B}(\hat{\lambda}) = \frac{3}{16n\alpha\lambda}, \quad (13)$$

Using Equations (12) and (13), we define the bias-corrected (BC) estimator as

$$\hat{\alpha}_{\text{BC}} = \hat{\alpha} - \hat{\mathcal{B}}(\hat{\alpha}), \quad \hat{\lambda}_{\text{BC}} = \hat{\lambda} - \hat{\mathcal{B}}(\hat{\lambda}). \quad (14)$$

Note that $\hat{\alpha}_{\text{BC}}$ and $\hat{\lambda}_{\text{BC}}$ defined in Equation (14) have bias of order $\mathcal{O}(n^{-2})$ as indicated in (11). Thus, it is expected that they have superior sampling properties relative to $\hat{\alpha}$ and $\hat{\lambda}$. We also empathize that the bias-corrected ML estimator for β in the original parametrization Equation (2) is obtained from $\hat{\beta} - (3\sqrt{\hat{\alpha}}\hat{\beta} + 1.5\sqrt{\hat{\beta}})/\sqrt{\hat{\alpha}n}$.

3.2 PARAMETRIC BOOTSTRAP BIAS CORRECTION

An alternative approach to analytically bias-corrected ML estimators is based on bootstrap resampling scheme (Efron and Tibshirani, 1993; Davison and Hinkley, 1997). In this method the bias correction is performed numerically without deriving analytical expression for the bias function. In fact, the parametric bootstrap bias correction (PB) estimates use the ML estimates of the data to generate pseudo-random samples from the distribution to estimate the bias and then subtract the bias from the ML estimates.

Let $\hat{\boldsymbol{\theta}}_{(\cdot)}$ be the average value of the ML estimator from B bootstrap replications, based on a pseudo-sample of size n generated from Equation (3) using the parameters of the ML estimates $\hat{\boldsymbol{\theta}}$. The estimated bias of $\hat{\boldsymbol{\theta}}$ is defined as $\hat{\mathcal{B}}(\hat{\boldsymbol{\theta}}) = \hat{\boldsymbol{\theta}}_{(\cdot)} - \hat{\boldsymbol{\theta}}$. Then, the bootstrap bias-corrected estimator is $\hat{\boldsymbol{\theta}}_{\text{PB}} = 2\hat{\boldsymbol{\theta}} - \hat{\boldsymbol{\theta}}_{(\cdot)}$.

4. NUMERICAL EVALUATIONS

In this section, we carry out a MC simulation study to compare the ML estimators and their bias-corrected versions. In addition, we illustrate the applicability of the CA distribution for bias corrections to the wind speed data.

4.1 SIMULATION STUDY

Our MC simulation study is conducted to compare the finite-sample behavior of the ML estimators and their bias-corrections obtained by Cox-Snell methodology (BC) and parametric bootstrap scheme (PB) for the parameters that index the CA distribution. For this purpose, we generate samples of size $n = 10, 20, 30, 40$ and 50 from Equation (3) considering $\alpha = 0.5, 1.0, 1.5, 2.0$ and 4.0 and fixed $\lambda = 1$, since it is a location parameter and the estimators are scale invariant. The behavior of PDF and HRF for these parameters values were illustrated in Figure 1. It is important to note that the mean, variance, skewness and kurtosis of a CA distributed random variable decrease as α increases.

To simulate random variables from a CA distribution, we generated samples from a random variable Y with inverse Gaussian distribution and we used the transformation $X = 1/\sqrt{Y}$.

To assess the performance of the methods under consideration, we calculated the bias and root mean-squared error (RMSE). The number of MC simulations was fixed at $M = 10,000$ and $B = 1,000$ bootstrap replicates were used. All simulations were carried out in Ox Console which is a matrix programming language with object-oriented support developed by Jurgen Doornik (Doornik, 2007).

Table 1 depicts the estimated bias and root mean-squared error, in parentheses, for different values of α and $\lambda = 1$. We can observe that all the estimates show the property of consistency, that is, the RMSEs decrease as sample size increases. We also note that the ML estimates of α are highly biased, particularly when the sample size is small. For instance, the biases of the ML estimates of α for $(n, \alpha) = (10, 0.5)$ and $(n, \alpha) = (10, 4)$ are approximately 22% and 169%, respectively. Also the biases of the ML estimates of α for $(n, \alpha) = (20, 0.5)$ and $(n, \alpha) = (20, 4)$ are approximately 9% and 68%, respectively. The estimates $\hat{\alpha}_{BC}$ and $\hat{\alpha}_{PB}$ clearly outperform the ML estimates as far as the bias goes. For example, the biases of the BC estimates of α for $(n, \alpha) = (10, 0.5)$ and $(n, \alpha) = (10, 4)$ are approximately 0.3% and 1.5%, respectively. The biases of the PB estimates of α for $(n, \alpha) = (10, 0.5)$ and $(n, \alpha) = (10, 4)$ are approximately 8.9% and 74.6%, respectively. Thus, the proposed estimators achieve substantial bias reduction, especially for the small and moderate sample sizes and therefore, we consider them as better alternatives of the ML estimates of α . We also observe that the bias-corrected estimates are closer to the true parameter values than the unadjusted estimates as sample size increases. Additionally, the estimated root mean-squared errors for α of the bias corrected estimates are smaller than those of the uncorrected estimates. On the other hand, the RMSE of λ are very similar for all estimators.

Now, in order to evaluate the overall performance of each estimation method with respect to the bias and root mean squared error, for each value of n , we use two measures introduced by Cribari-Neto and Vasconcellos (2002). The authors called these quantities as integrated bias squared norm and average root mean squared error. They are calculated as follows

$$\text{IBSQ}_{(k)} = \sqrt{\frac{1}{16} \sum_{h=1}^{16} (r_{h,k})^2}, \quad \text{ARMSE}_{(k)} = \frac{1}{16} \sum_{h=1}^{16} \text{RMSE}_{h,k},$$

Table 1. Estimated bias (root mean-squared error) for α and λ , ($\lambda = 1.0$).

α	n	Estimator of α			Estimator of λ		
		ML	BC	PB	ML	BC	PB
0.5	10	0.2191 (0.5098)	0.0034 (0.3223)	-0.0891 (0.2778)	0.0351 (0.1646)	0.0018 (0.1667)	0.0029 (0.1662)
	20	0.0908 (0.2345)	0.0022 (0.1838)	-0.0135 (0.1787)	0.0173 (0.1142)	-0.0003 (0.1149)	-0.0000 (0.1149)
	30	0.0571 (0.1686)	0.0014 (0.1428)	-0.0048 (0.1411)	0.0111 (0.0924)	-0.0009 (0.0929)	-0.0008 (0.0929)
	40	0.0410 (0.1369)	0.0004 (0.1208)	-0.0029 (0.1200)	0.0082 (0.0798)	-0.0009 (0.0801)	-0.0008 (0.0801)
	50	0.0324 (0.1174)	0.0005 (0.1061)	-0.0016 (0.1057)	0.0063 (0.0715)	-0.0010 (0.0717)	-0.0010 (0.0717)
1.0	10	0.4330 (0.9830)	0.0031 (0.6178)	-0.1810 (0.5368)	0.0171 (0.1131)	0.0004 (0.1138)	0.0006 (0.1137)
	20	0.1775 (0.4620)	0.0009 (0.3625)	-0.0303 (0.3526)	0.0085 (0.0793)	-0.0004 (0.0795)	-0.0003 (0.0795)
	30	0.1149 (0.3370)	0.0034 (0.2851)	-0.0090 (0.2818)	0.0059 (0.0648)	-0.0001 (0.0649)	-0.0001 (0.0650)
	40	0.0844 (0.2722)	0.0031 (0.2394)	-0.0035 (0.2380)	0.0046 (0.0559)	0.0000 (0.0560)	0.0000 (0.0560)
	50	0.0667 (0.2334)	0.0027 (0.2103)	-0.0015 (0.2094)	0.0034 (0.0501)	-0.0003 (0.0501)	-0.0003 (0.0501)
1.5	10	0.6543 (1.5198)	0.0080 (0.9603)	-0.2688 (0.8293)	0.0121 (0.0927)	0.0010 (0.0931)	0.0011 (0.0930)
	20	0.2657 (0.6906)	0.0008 (0.5419)	-0.0459 (0.5272)	0.0064 (0.0653)	0.0005 (0.0654)	0.0005 (0.0654)
	30	0.1653 (0.4970)	-0.0013 (0.4218)	-0.0197 (0.4173)	0.0040 (0.0530)	-0.0000 (0.0530)	-0.0000 (0.0530)
	40	0.1213 (0.4035)	-0.0003 (0.3559)	-0.0100 (0.3539)	0.0027 (0.0457)	-0.0004 (0.0458)	-0.0003 (0.0458)
	50	0.0963 (0.3489)	0.0005 (0.3152)	-0.0056 (0.3140)	0.0022 (0.0410)	-0.0002 (0.0410)	-0.0002 (0.0410)
2.0	10	0.8784 (2.0228)	0.0149 (1.2756)	-0.3550 (1.1005)	0.0090 (0.0796)	0.0007 (0.0798)	0.0007 (0.0798)
	20	0.3588 (0.9360)	0.0050 (0.7348)	-0.0575 (0.7144)	0.0042 (0.0559)	-0.0003 (0.0560)	-0.0003 (0.0560)
	30	0.2319 (0.6808)	0.0087 (0.5762)	-0.0161 (0.5694)	0.0026 (0.0456)	-0.0004 (0.0457)	-0.0004 (0.0457)
	40	0.1707 (0.5483)	0.0079 (0.4820)	-0.0054 (0.4788)	0.0020 (0.0397)	-0.0003 (0.0397)	-0.0003 (0.0397)
	50	0.1317 (0.4702)	0.0038 (0.4244)	-0.0044 (0.4227)	0.0015 (0.0354)	-0.0003 (0.0354)	-0.0003 (0.0354)
4.0	10	1.6928 (3.8947)	-0.0150 (2.4553)	-0.7463 (2.1403)	0.0053 (0.0565)	0.0011 (0.0566)	0.0011 (0.0566)
	20	0.6965 (1.8473)	-0.0080 (1.4543)	-0.1329 (1.4157)	0.0027 (0.0398)	0.0004 (0.0398)	0.0004 (0.0398)
	30	0.4358 (1.3260)	-0.0078 (1.1271)	-0.0573 (1.1144)	0.0018 (0.0325)	0.0003 (0.0325)	0.0003 (0.0325)
	40	0.3110 (1.0752)	-0.0123 (0.9521)	-0.0385 (0.9465)	0.0014 (0.0282)	0.0003 (0.0282)	0.0003 (0.0282)
	50	0.2406 (0.9169)	-0.0138 (0.8318)	-0.0301 (0.8289)	0.0011 (0.0253)	0.0002 (0.0253)	0.0002 (0.0253)

where $r_{h,k}$ and $\text{RMSE}_{h,k}$ correspond to the 16 different values of the bias and the root mean squared errors of each estimator given in Table 1. The results are reported in Tables 2-3.

From Table 2, we see that integrated bias squared norm of the corrected estimates (BC and PB) are smaller than ML estimates for both parameter α and λ . From Table 2, we can see that the average root mean-squared error of the corrected estimates (BC and PB) are smaller than ML estimates for α , while for λ the ARMSE are quite similar. Therefore, these simulation results show that second-order bias reduction is quite successful in bringing the corrected estimates closer to their true values.

Table 2. Integrated bias squared norm.

n	Estimator of α			Estimator of λ		
	ML	BC	PB	ML	BC	PB
10	0.9274	0.0103	0.3990	0.0189	0.0011	0.0015
20	0.3806	0.0043	0.0695	0.0094	0.0004	0.0004
30	0.2398	0.0055	0.0284	0.0061	0.0005	0.0004
40	0.1728	0.0067	0.0181	0.0045	0.0005	0.0004
50	0.1342	0.0065	0.0139	0.0035	0.0005	0.0005

Table 3. Average root mean-squared error.

n	Estimator of α			Estimator of λ		
	ML	BC	PB	ML	BC	PB
10	2.1352	1.3464	1.1701	0.1077	0.1086	0.1084
20	1.0034	0.7892	0.7680	0.0752	0.0755	0.0755
30	0.7226	0.6135	0.6066	0.0611	0.0613	0.0613
40	0.5852	0.5172	0.5141	0.0528	0.0530	0.0530
50	0.5004	0.4532	0.4515	0.0473	0.0474	0.0474

4.2 WIND REAL DATA MODELING

The data consist of annual maximum wind speed of six weather stations localized in state of Tocantins, Brazil. The data were obtained from the website <http://www.inmet.gov.br/portal>. Some descriptive statistics of the observed annual maximum wind speed for the stations are summarized in Table 4. Note that the values of skewness are positive for four stations, indicating that the data are right-skewed.

Table 4. Descriptive statistics of the wind speed data for all weather stations.

Station	Period	n	Min	Mean	Med	Max	SD	Skewn	Kurt
82659	1980-2016	34	1.4667	3.0441	3.0667	5.0000	1.1491	0.1541	1.5672
82863	1977-2016	40	2.5667	4.8862	4.6667	8.3333	1.6084	0.4240	2.0874
83033	1993-2016	21	3.6667	5.4817	5.4667	7.1000	1.0460	-0.1572	2.1114
83064	1961-2016	56	2.2667	3.9939	3.9750	6.0000	0.9736	0.1973	2.3523
83228	1975-2016	42	3.1000	4.3657	4.3333	5.6667	0.6096	-0.1140	2.6386
83235	1961-2016	56	2.3333	3.9994	3.6667	6.6667	0.9424	0.8900	3.4780

In Table 5 we report the ML estimates and the bias corrections estimates along with the asymptotic standard errors calculated from Equation (9). We can observe that the maximum likelihood estimates of α and λ are greater than the second order bias corrected estimates for all stations, this suggests that the ML estimates are overestimating the true value of the parameters. We also observe that the corrected ML estimates of α have smaller standard errors than the uncorrected estimates.

In order to test whether the data sets fits the CA distribution and whether the bias-corrected estimates yield better fits than the uncorrected estimates, we perform the goodness-of-fit tests based on Kolmogorov-Smirnov (KS), Cramér-von-Mises (CM) and Anderson-Darling (AD) statistics. The p -values of these statistics are shown in Table 8. We have used the function *mledist* from *fitdistrplus* library, (Delignette-Muller, 2015), available in *R* environment, (R Core Team, 2017), to find the ML estimates. The p -values associated with Kolmogorov-Smirnov, Anderson-Darling and Cramér-von Mises tests were calculated using 10,000 nonparametric bootstrap resamples applying the functions *ks.test*, *ad.test* and *cvm.test*, available in *gofTest* (Faraway et al., 2014) *R* library. Here we are to note that CA distribution can be used to model the annual maximum wind speed data for the six stations. We can see from Table 8 that the p -values of KS, CM and AD computed from bias-corrected estimators are greater than the uncorrected estimator (except KS for one station). This means that bias corrected estimates provide better fits than the ML estimates. This conclusion is also supported by the empirical and fitted CDF plots in Figure 2. Furthermore, in Table 6, we compare the suitability of the CA distribution against eight commonly used probability distributions to modeling wind speed data. The assessment of the goodness-of-fit is based on the log-likelihood values, since all distributions have the same number of parameters. The results are reported in Table 6 and we can see that CA distribution is the best model among the others. The superscripts indicates the rank obtained by the estimation method (the smaller the better). The line named as rank total (TR) shows the sum of the ranks.

Table 5. Point estimates (standard errors) for all weather stations.

Station	Estimator of α			Estimator of λ		
	ML	BC	PB	ML	BC	PB
82659	0.1081 (0.0262)	0.0986 (0.0239)	0.0978 (0.0237)	2.4341 (0.1844)	2.4132 (0.1931)	2.4121 (0.1939)
82863	0.0559 (0.0125)	0.0517 (0.0116)	0.0515 (0.0115)	4.1773 (0.2365)	4.1572 (0.2459)	4.1627 (0.2464)
83033	0.1148 (0.0354)	0.0984 (0.0304)	0.0962 (0.0297)	5.1707 (0.2277)	5.1556 (0.2459)	5.1539 (0.2487)
83064	0.1359 (0.0257)	0.1286 (0.0243)	0.1279 (0.0242)	3.6335 (0.1282)	3.6268 (0.1317)	3.6250 (0.1321)
83228	0.3323 (0.0725)	0.3086 (0.0673)	0.3075 (0.0671)	4.2329 (0.0946)	4.2297 (0.0982)	4.2315 (0.0984)
83235	0.1639 (0.0310)	0.1551 (0.0293)	0.1540 (0.0291)	3.7170 (0.1167)	3.7115 (0.1200)	3.7098 (0.1204)




Figure 2. Empirical and fitted CDFs for all examined stations.

Table 6. Negative of the log likelihood values, $-2\log(L)$, of the competing distributions.

Station	CA	Weibull	Gamma	Log-normal	Log-logistic	IG	Gumbel	BS	Nakagami
82659	101.4117 ¹	103.0848 ⁵	103.1692 ⁶	103.6138 ⁷	107.4095 ⁹	103.0738 ⁴	104.3379 ⁸	102.9892 ²	103.0680 ³
82863	145.7202 ¹	149.6565 ⁸	147.2638 ⁵	146.9742 ⁴	150.4022 ⁹	146.5940 ³	147.4252 ⁶	146.5772 ²	148.0746 ⁷
83033	61.3816 ⁴	60.0757 ¹	61.0546 ³	61.6314 ⁷	62.5727 ⁸	61.6019 ⁶	63.3180 ⁹	61.5889 ⁵	60.6320 ²
83064	154.2422 ³	155.7540 ⁷	154.1821 ²	155.1365 ⁶	157.5565 ⁹	155.0057 ⁵	157.1036 ⁸	154.9604 ⁴	153.9516 ¹
83228	78.1260 ⁴	77.5426 ³	77.4511 ²	78.2403 ⁷	79.0705 ⁸	78.2372 ⁶	83.3099 ⁹	78.2280 ⁵	76.8940 ¹
83235	143.7640 ⁵	156.2722 ⁹	145.2559 ⁷	143.5298 ⁴	144.2417 ⁶	143.4797 ²	142.3888 ¹	143.5114 ³	147.7691 ⁸
RT	18 ¹	33 ⁶	25 ⁴	35 ⁷	49 ⁹	26 ⁵	41 ⁸	21 ²	22 ³

Table 7. Young test (p -values) comparing CA distribution with others.

Weibull	Gamma	Log-normal	Log-logistic	IG	Gumbel	BS	Nakagami
0.595 (0.276)	2.146 (0.016)	3.782 (0.000)	8.895 (0.000)	2.538 (0.006)	5.876 (0.000)	2.773 (0.003)	0.882 (0.189)
1.522 (0.064)	2.171 (0.015)	2.460 (0.007)	4.628 (0.000)	1.806 (0.035)	2.156 (0.016)	1.959 (0.025)	1.479 (0.070)
-0.650 (0.742)	-0.788 (0.785)	1.378 (0.084)	1.218 (0.112)	1.564 (0.059)	2.545 (0.005)	1.482 (0.069)	-1.014 (0.845)
0.497 (0.310)	-0.072 (0.529)	2.107 (0.018)	1.929 (0.027)	2.267 (0.012)	3.379 (0.000)	2.203 (0.014)	-0.192 (0.576)
-0.168 (0.567)	-1.282 (0.900)	0.872 (0.192)	0.654 (0.256)	1.090 (0.138)	2.854 (0.002)	1.008 (0.157)	-1.233 (0.891)
3.457 (0.000)	1.943 (0.026)	-0.555 (0.711)	0.244 (0.404)	-0.799 (0.788)	-0.686 (0.754)	-0.750 (0.773)	2.375 (0.009)

Table 8. p -values associated to goodness-of-fit measures for all weather stations.

Station	KS			CM			AD		
	ML	BC	PB	ML	BC	PB	ML	BC	PB
82659	0.4697	0.5185	0.5205	0.3129	0.3970	0.4039	0.3247	0.4056	0.4112
82863	0.6574	0.7285	0.7192	0.7725	0.8539	0.8515	0.7825	0.8570	0.8547
83033	0.9390	0.9763	0.9775	0.8936	0.9252	0.9262	0.8599	0.9042	0.9047
83064	0.8883	0.8736	0.8684	0.7438	0.7689	0.7677	0.7557	0.7962	0.7976
83228	0.4790	0.4961	0.5090	0.6607	0.6748	0.6836	0.7368	0.7534	0.7578
83235	0.2996	0.3054	0.3097	0.4005	0.3801	0.3795	0.4824	0.4735	0.4731

Therefore, using the interpretation for λ given in Section 1, the estimates given in Table 5 can be interpreted as follows. The most frequent wind speed at: station 82659 is around 2.4; station 82863 is around 4.2; station 83033 is around 5.2; station 83064 is around 3.6; station 83228 is around 4.2; station 83235 is around 3.7.

5. CONCLUSIONS

In this paper, we have adopted a corrective approach to derive analytical expressions for the second order biases of the maximum likelihood estimators of the parameters of the Chaudhry-Ahmad distribution. Furthermore, we have also considered an alternative bias-correction mechanism through bootstrap resampling. The biases of the proposed estimators are of order $\mathcal{O}(n^{-2})$, whereas for the maximum likelihood estimators they are of order $\mathcal{O}(n^{-1})$, indicating that the proposed estimates converge to their true value considerably faster than those of the maximum likelihood estimates.

The numerical evidence shows that the proposed bias corrected estimators are quite attractive because they outperform the maximum likelihood estimates in terms of biases, integrated bias squared norm and root mean-squared error. Further, our analytic bias correction is found to be superior to the alternative of bias-correction via the bootstrap in terms of bias reduction. The proposed bias-corrected estimators are strongly recommended over maximum likelihood estimator, especially when the sample size is small or moderate since it has smaller bias and root mean-squared error

ACKNOWLEDGEMENTS

The authors are thankful to the referees and to the Editors-in-Chiefs for many valuable suggestions. Josmar Mazucheli gratefully acknowledge the partial financial support from Fundação Araucária (Grant 064/2019 - UEM/Fundação Araucária).

REFERENCES

- Abramowitz, M. and Stegun, I.A., 1974. Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables. Dover, New York.
- Chaudhry, M.A. and Ahmad, M., 1993. On a probability function useful in size modelling. *Canadian Journal of Forest Research*, 23, 1679–1683.
- Cox, D.R. and Hinkley, D.V., 1979. Theoretical Statistics. CRC Press, Boca Raton, FL, US.
- Cox, D.R. and Reid, N., 1987. Parameter orthogonality and approximate conditional inference. *Journal of the Royal Statistical Society B*, 49, 1–39.
- Cox, D.R. and Snell, E.J., 1968. A general definition of residuals. *Journal of the Royal Statistical Society B*, 30, 248–275.
- Cribari-Neto, F. and Vasconcellos, K.L.P., 2002. Nearly unbiased maximum likelihood estimation for the beta distribution. *Journal of Statistical Computation and Simulation*, 72, 107–118.
- Davison, A.C. and Hinkley, D.V., 1997. Bootstrap Methods and Their Applications. Cambridge University Press, Cambridge.
- Delignette-Muller, M.L. and Dutang, C., 2015. fitdistrplus: An R package for fitting distributions. *Journal of Statistical Software*, 64, 1–34.
- Doornik, J.A., 2007. Object-Oriented Matrix Programming Using Ox. Timberlake Consultants Press and Oxford, London.
- Edwards, A.W.F., 1992. Likelihood. Johns Hopkins University Press, Baltimore.
- Efron, B., 1982. The Jackknife, the Bootstrap and other Resampling Plans. SIAM.
- Efron, B. and Tibshirani, R.J., 1993. An Introduction to the Bootstrap. Chapman and Hall, New York.
- Faraway, J., Marsaglia, G., Marsaglia, J., and Baddeley, A., 2014. goftest: Classical goodness-of-fit tests for univariate distributions. R package version 1.0-2.
- Giles, D.E., 2012a. Bias reduction for the maximum likelihood estimators of the parameters in the Half-Logistic distribution. *Communication in Statistics: Theory and Methods*, 41, 212–222.
- Giles, D.E., 2012b. A note on improved estimation for the Topp-Leone distribution. Technical Report. Department of Economics, University of Victoria.
- Giles, D.E. and Feng, H., 2009. Bias of the maximum likelihood estimators of the two-parameter Gamma distribution revisited. Technical Report. Department of Economics, University of Victoria.
- Giles, D.E., Feng, H., and Godwin, R.T., 2013. On the bias of the maximum likelihood estimator for the two-parameter Lomax distribution. *Communications in Statistics: Theory and Methods*, 42, 1934–1950.
- Koutras, V.P., 2011. Two-level software rejuvenation model with increasing failure rate degradation. *Dependable Computer Systems*, 97, 101–115.
- Lagos-Álvarez, B., Jiménez-Gamero, M.D., and Alba-Fernández, V., 2011. Bias correction in the type I generalized Logistic distribution. *Communications in Statistics: Simulation and Computation*, 40, 511–531.

- Lai, M.T., 2013. Optimum number of minimal repairs for a system under increasing failure rate shock model with cumulative repair-cost limit. *International Journal of Reliability and Safety*, 7, 95–107.
- Lehmann, E.J. and Casella, G., 1998. *Theory of Point Estimation*. Springer, New York.
- Lemonte, A.J., Cribari-Neto, F., and Vasconcellos, K.L., 2007. Improved statistical inference for the two-parameter Birnbaum-Saunders distribution. *Computational Statistics and Data Analysis*, 51, 4656 – 4681.
- Ling, X. and Giles, D.E., 2014. Bias reduction for the maximum likelihood estimator of the parameters of the generalized Rayleigh family of distributions. *Communications in Statistics: Theory and Methods*, 43, 1778–1792.
- Mazucheli, J., Bapat, S.R., and Menezes, A.F.B., 2020. A new one-parameter unit-Lindley distribution. *Chilean Journal of Statistics*, 11, 53–67.
- Mazucheli, J., and Dey, S., 2018. Bias-corrected maximum likelihood estimation of the parameters of the generalized half-normal distribution. *Journal of Statistical Computation and Simulation*, 88, 1027–1038.
- Mazucheli, J., Menezes, A.F.B., and Dey, S., 2018. Improved maximum likelihood estimators for the parameters of the unit-gamma distribution. *Communications in Statistics: Theory and Methods*, 47, 3767–3778.
- Nanos, N. and Montero, G., 2001. *Spatial Prediction of Diameter Distribution Models in Forestry*. Springer, Dordrecht.
- Nanos, N. Tadesse, W., Montero, G., Gil, L., and Alia, R., 2000. Modelling resin production distributions for pinus pinaster by using two probability functions. *Annals of Forest Science*, 57, 369–377.
- Pearson, K., 1893. Contributions to the mathematical theory of evolution. *Proceedings of the Royal Society of London*, 54, 329–333.
- Pearson, K., 1895. Contributions to the mathematical theory of evolution, II: Skew variation in homogeneous material. *Philosophical Transactions of the Royal Society of London A*, A186, 343–414.
- Pearson, K., 1901. Mathematical contributions to the theory of evolution, X: Supplement to a memoir on skew of variation. *Philosophical Transactions of the Royal Society of London A*, 197, 443–459.
- Pearson, K., 1916. Mathematical contributions to the theory of evolution, XIX: Second supplement to a memoir on skew of variation. *Philosophical Transactions of the Royal Society of London A*, 216, 429–457.
- R Core Team, 2017. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. Vienna, Austria.
- Reath, J., 2016. Improved parameter estimation of the Log-Logistic distribution with applications. Ph.D. thesis. Michigan Technological University.
- Saha, K. and Paul, S., 2005. Bias-corrected maximum likelihood estimator of the negative Binomial dispersion parameter. *Biometrics*, 61, 179–185.
- Sankaran, P.G., Nair, N.U., and Sindhu, T.K., 2003. A generalized Pearson system useful in reliability analysis. *Statistical Papers*, 44, 125–130.
- Schwartz, J. and Giles, D.E., 2016. Bias-reduced maximum likelihood estimation of the zero-inflated Poisson distribution. *Communications in Statistics: Theory and Methods*, 45, 465–478.
- Schwartz, J., Godwin, R.T., and Giles, D.E., 2013. Improved maximum-likelihood estimation of the shape parameter in the Nakagami distribution. *Journal of Statistical Computation and Simulation*, 83, 434–445.
- Shakil, M., Kibria, B.M.G., and Singh, J.N., 2010. A new family of distributions based on the generalized Pearson differential equation with some applications. *Austrian Journal of Statistics*, 39, 259–278.

- Shakil, M., Kibria, B.M.G., and Singh, J.N., 2016. Review on generalized Pearson system of probability distributions. *Journal of Mathematical Sciences and Mathematics Education*, 11, 13–33.
- Stavroyiannis, S., 2014. On the generalised Pearson distribution for application in financial time series modelling. *Global Business and Economics Review*, 16, 1–14.
- Teimouri, M. and Nadarajah, S., 2013. Bias corrected MLEs for the Weibull distribution based on records. *Statistical Methodology*, 13, 12–24.
- Teimouri, M. and Nadarajah, S., 2016. Bias corrected MLEs under progressive type-II censoring scheme. *Journal of Statistical Computation and Simulation*, 86, 2714–2726.
- Tsarouhas, P.H. and Arvanitoyannis, I.S., 2010. Reliability and maintainability analysis of bread production line. *Critical Reviews in Food Science and Nutrition*, 50, 327–343.
- Wang, M. and Wang, W., 2017. Bias-corrected maximum likelihood estimation of the parameters of the weighted Lindley distribution. *Communications in Statistics: Theory and Methods*, 46, 530–545.
- Woosley, R.L., Cossman, J., 2007. Drug development and the FDA’s critical path initiative. *Public Policy*, 81, 129–133.
- Zhang, G. and Liu, R., 2015. Bias-corrected estimators of scalar skew Normal. *Communications in Statistics: Simulation and Computation*, 46, 831–839.

INFORMATION FOR AUTHORS

The editorial board of the Chilean Journal of Statistics (ChJS) is seeking papers, which will be refereed. We encourage the authors to submit a PDF electronic version of the manuscript in a free format to the Editors-in-Chief of the ChJS (E-mail: chilean.journal.of.statistics@gmail.com). Submitted manuscripts must be written in English and contain the name and affiliation of each author followed by a leading abstract and keywords. The authors must include a "cover letter" presenting their manuscript and mentioning: "We confirm that this manuscript has been read and approved by all named authors. In addition, we declare that the manuscript is original and it is not being published or submitted for publication elsewhere".

PREPARATION OF ACCEPTED MANUSCRIPTS

Manuscripts accepted in the ChJS must be prepared in Latex using the ChJS format. The Latex template and ChJS class files for preparation of accepted manuscripts are available at <http://chjs.mat.utfsm.cl/files/ChJS.zip>. Such as its submitted version, manuscripts accepted in the ChJS must be written in English and contain the name and affiliation of each author, followed by a leading abstract and keywords, but now mathematics subject classification (primary and secondary) are required. AMS classification is available at <http://www.ams.org/mathscinet/msc/>. Sections must be numbered 1, 2, etc., where Section 1 is the introduction part. References must be collected at the end of the manuscript in alphabetical order as in the following examples:

Arellano-Valle, R., 1994. Elliptical Distributions: Properties, Inference and Applications in Regression Models. Unpublished Ph.D. Thesis. Department of Statistics, University of São Paulo, Brazil.

Cook, R.D., 1997. Local influence. In Kotz, S., Read, C.B., and Banks, D.L. (Eds.), Encyclopedia of Statistical Sciences, Vol. 1., Wiley, New York, pp. 380-385.

Rukhin, A.L., 2009. Identities for negative moments of quadratic forms in normal variables. Statistics and Probability Letters, 79, 1004-1007.

Stein, M.L., 1999. Statistical Interpolation of Spatial Data: Some Theory for Kriging. Springer, New York.

Tsay, R.S., Peña, D., and Pankratz, A.E., 2000. Outliers in multivariate time series. Biometrika, 87, 789-804.

References in the text must be given by the author's name and year of publication, e.g., Gelfand and Smith (1990). In the case of more than two authors, the citation must be written as Tsay et al. (2000).

COPYRIGHT

Authors who publish their articles in the ChJS automatically transfer their copyright to the Chilean Statistical Society. This enables full copyright protection and wide dissemination of the articles and the journal in any format. The ChJS grants permission to use figures, tables and brief extracts from its collection of articles in scientific and educational works, in which case the source that provides these issues (Chilean Journal of Statistics) must be clearly acknowledged.